

THE SEIBERG-WITTEN MAP IN NONCOMMUTATIVE FIELD THEORY: AN ALTERNATIVE INTERPRETATION

Subir Ghosh

Physics and Applied Mathematics Unit,
Indian Statistical Institute,
203 B. T. Road, Calcutta 700108,
India.

Abstract:

In this article, an alternative interpretation of the Seiberg-Witten map in non-commutative field theory is provided. We show that the Seiberg-Witten map can be induced in a geometric way, by a field dependent co-ordinate transformation that connects noncommutative and ordinary space-times. Furthermore, in continuation of our earlier works, it has been demonstrated here that the above (field dependent co-ordinate) transformations are present in a gauge fixed version of the relativistic spinning particle model, embedded in the Batalin-Tyutin extended space. We emphasize that the space-time non-commutativity emerges naturally from the particle *spin* degrees of freedom. Contrary to similarly motivated works, the non-commutativity is not imposed here in an *ad-hoc* manner.

Keywords: Seiberg-Witten map, non-commutative space-time, spinning particle.

PACS Numbers: 02.40.Gh; 11.10.Ef; 11.90.+t

INTRODUCTION

Historically the Non-Commutative (NC) spacetime was introduced by Snyder [1] as a regularization to tame the short distance singularities, inherent in a Quantum Field Theory (QFT). In some sense, this extra structure might appear as a natural generalization of the phase space non-commutativity in quantum mechanics. The softening of the divergence is obviously because NC in spacetime can introduce a lower bound in the continuity of spacetime, just as $\hbar$ does in the quantum phase space. The advantage of NC as a regularization is that the computational scheme requires very little changes from the ordinary spacetime and in some forms of NC [1], (for more recent works see [2, 3, 4]), manifest Lorentz invariance can be maintained. ¹All the same, the idea of Snyder [1] lost popularity due to the success of renormalization techniques in QFT. Also, now we know [6] that noncommutativity is not of much use as a regulating scheme.

However, in more recent times, NCQFT has matured as an area of intense research activity [6]. It has been established by Seiberg and Witten [7] that the existence of non-commutativity in (open) string boundaries in the presence of a constant two-form Neveu-Schwarz field results in NC D-branes to which the open string endpoints are attached. (Hamiltonian formulation of the above open string boundary noncommutativity is discussed in [8].) This renders QFTs living on the D-brane noncommutative. The authors in [7] also propose an explicit mapping between NC and ordinary space-time dynamical variables. This celebrated map goes by the name of the Seiberg-Witten Map (SWM) [7]. Actually the SWM is the culmination of a much deeper understanding between the connection of QFTs in NC and ordinary space-time. It has been shown in [7] that the appearance of NCQFT is dependent on the choice of regularization and in fact a QFT in ordinary spacetime and an NCQFT both can describe the same underlying QFT. At least to the lowest non-trivial order in $\theta_{\mu\nu}$ ², the non-commutativity parameter,

$$[x_\mu, x_\nu] = i\theta_{\mu\nu}, \tag{1}$$

the SWM can be exploited to convert a NCQFT to its counterpart living in ordinary space-time, in which the effects of non-commutativity appear as local interaction terms, supplemented by $\theta_{\mu\nu}$. In the more popular form of NCQFT, $\theta_{\mu\nu}$ is taken to be constant. This can lead to very striking signatures in particle physics phenomenology in the form of Lorentz symmetry breakdown, new interaction vertices etc. [10].

The SWM plays a pivotal role in our understanding of the NCQFT by directly making contact between NCQFT and QFT in ordinary space-time. This has led to new results in the form of axial anomaly [11] in an NC interacting theory of fermions coupled to gauge fields. The previous results [12] are incomplete as they do not conform to the SWM. Furthermore, the SWM is crucial in generalizing [13] to NC space-time, the duality between Maxwell-Chern-Simons and Self-Dual theories in 2+1-dimensions [14]. In the context of bosonization of the massive Thirring model in 2+1-dimensions [15], SWM reveals that the resulting theory is different from the NC self-dual model [16], in contrast to ordinary space-time.

¹The ideas of [3] is discussed in a field theoretic setting in [5].

²The SWM for higher orders in θ are discussed in [9].

However, as it stands, the SWM is linked exclusively to NC *gauge* theory, since the original derivation of the SWM [7] hinges on the concept of identifying gauge orbits in NC and ordinary space-times. In the explicit form of the SWM [7], the non-commutativity of the *space-time* in which the NC gauge field lives, is not manifest at all since the map is a relation between the NC and ordinary gauge fields and gauge transformation parameters, all having ordinary space-time coordinates as their arguments.

On the other hand, possibly it would have been more natural to consider first a map between NC and ordinary space-time and subsequently to induce the SWM through the change in the space-time argument of the gauge field from the ordinary to NC one. Precisely this type of a geometrical reformulation of the SWM is proposed in the present work.

The paper is organized as follows: **Section II** describes the motivation and an outline of our ideas. The SWM is analysed in a different light in **Section III**. This constitutes our alternative interpretation of the SWM. **Section IV** is devoted to a discussion on the Batalin-Tyutin extension and the ordinary and noncommutative spacetime mapping. The Batalin-Tyutin extension and SWM are bridged through the process of gauge fixing in **Section V**. The paper ends with a conclusion and future prospects in **Section VI**.

II. MOTIVATION AND AN OUTLINE OF THE PRESENT WORK

In the canonical quantization prescription, the Poisson Bracket algebra is elevated to quantum commutator algebra by the replacement

$$\{A, B\} \rightarrow \frac{1}{i}[\hat{A}, \hat{B}].$$

But presence of *constraints* may demand a modification in the Poisson Bracket algebra, leading to the Dirac Bracket algebra [17], which are subsequently identified to the commutators,

$$\{A, B\}_{DB} \rightarrow \frac{1}{i}[\hat{A}, \hat{B}].$$

However, complications in this formalism can arise, (in particular in case of non-linear constraints), where the Dirac Bracket algebra itself becomes operator valued. To overcome this, Batalin and Tyutin [18] have developed a systematic scheme in which all the physical variables are mapped in an extended canonical phase space, consisting of auxiliary degrees of freedom besides the physical ones, with all of them enjoying canonical free Poisson Bracket algebra. In this formalism, the ambiguity of using (operator valued) Dirac Brackets as quantum commutators does not arise.

In the spinning particle model [19] the canonical $\{x_\mu, x_\nu\} = 0$ Poisson Bracket changes to an operator valued Dirac Bracket,

$$\{x_\mu, x_\nu\}_{DB} = -\frac{S_{\mu\nu}}{M^2}, \quad (2)$$

due to the presence of constraints. In the above, the dynamical variable $S_{\mu\nu}$ represents the spin angular momentum and M is the mass of the particle. This forces us to exploit the Batalin-Tyutin prescription [18].

In a recent paper [3], we have constructed a mapping of the form,

$$\{x_\mu, x_\nu\} = 0 \quad , \quad x_\mu \rightarrow \hat{x}_\mu \quad ; \quad \{\hat{x}_\mu, \hat{x}_\nu\} = \theta_{\mu\nu}, \quad (3)$$

which bridges the noncommutative and ordinary space-times. Note that $\hat{x}_\mu$ lives in the Batalin-Tyutin [18] extended space and is of the generic form $\hat{x}_\mu = x_\mu + X_\mu$, where X_μ consists of physical and auxiliary degrees of freedom. Explicit expressions for X_μ are to be found later [3].

This space-time map induces in a natural way the following map between noncommutative and ordinary degrees of freedom,

$$\lambda(x) \rightarrow \lambda(\hat{x}) \rightarrow \hat{\lambda}(x) \quad , \quad A_\mu(x) \rightarrow A_\mu(\hat{x}) \rightarrow \hat{A}_\mu(x). \quad (4)$$

Here $\hat{\lambda}$ and $\hat{A}_\mu$ are the NC counterparts of λ and A_μ , the abelian gauge transformation parameter and the gauge field respectively and $\hat{x}_\mu$ and x_μ are the NC and ordinary space-time co-ordinates.

On the other hand, there also exists the SWM [7] which interpolates between noncommutative and ordinary variables,

$$\lambda(x) \rightarrow \hat{\lambda}(x) \quad , \quad A_\mu(x) \rightarrow \hat{A}_\mu(x). \quad (5)$$

It is only logical that the above two schemes ((3-4) and 5) can be related. In the present work we have precisely done that. The formulation [3] (3-4) being the more general one, we have explicitly demonstrated how it can be reduced to the SWM [7], in a particular gauge. This incidentally demonstrates the correctness of the procedure. The above idea was hinted in [3].³

In this context, let us put the present work in its proper perspective. Recently a number of works have appeared with the motivation of recovering the SWM in a geometric way, without invoking the gauge theory principles [20]. However, the noncommutative feature of the space-time plays no direct role in the above mentioned re-derivations of the SWM, with non-commutativity just being postulated in an *ad hoc* way. In the present work, we have shown how one can construct a noncommutative sector inside an extended phase space, in a relativistically covariant way. More importantly, we have shown explicitly how this generalized map can be reduced to the SWM under certain approximations. Interestingly, this extended space is physically significant and well studied: It is the space of the relativistic spinning particle [3, 19]. Hence it might be intuitively appealing to think that the NC space-time is endowed with spin degrees of freedom, as compared to the ordinary configuration space, since the spin variables directly generate the NC [5]. The analogue of the gauge field is also identified inside this phase space, without any need to consider external fields. This situation is to be contrasted with the NC arising from the background magnetic field in the well known Landau problem [6] of a charge moving in a plane in the presence of a strong, perpendicular magnetic field.

³The present analysis being classical, (non)commutativity is to be interpreted in the sense of Poisson or Dirac Brackets.

III. SEIBERG-WITTEN MAP: AN ALTERNATIVE INTERPRETATION

The genesis of the SWM is the observation [7] that the non-commutativity in string theory depends on the choice of the regularization scheme: it appears in *e.g.* point-spitting regularization whereas it does not show up in Pauli Villars regularization. This feature, among other things, has prompted Seiberg and Witten [7] to suggest the map connecting the NC gauge fields and gauge transformation parameter to the ordinary gauge field and gauge transformation parameter. The explicit form of the SWM [7], for abelian gauge group, to the first non-trivial order in the NC parameter $\theta_{\mu\nu}$ is the following,

$$\begin{aligned}\hat{\lambda}(x) &= \lambda(x) + \frac{1}{2}\theta^{\mu\nu}A_\nu(x)\partial_\mu\lambda(x) , \\ \hat{A}_\mu(x) &= A_\mu(x) + \frac{1}{2}\theta^{\sigma\nu}A_\nu(x)F_{\sigma\mu}(x) + \frac{1}{2}\theta^{\sigma\nu}A_\nu(x)\partial_\sigma A_\mu(x).\end{aligned}\tag{6}$$

The above relation (6) is an $O(\theta)$ solution of the general map [7],

$$\hat{A}_\mu(A + \delta_\lambda A) = \hat{A}_\mu(A) + \hat{\delta}_\lambda \hat{A}_\mu(A),\tag{7}$$

which is based on identifying gauge orbits in NC and ordinary space-time.

First let us show that it is indeed possible to re-derive the SWM using geometric objects. We rewrite the SWM (6) in the following way,

$$\hat{\lambda}(x) = \lambda(x) + \frac{1}{2}\{\delta_f[\lambda(x)] - (\lambda(x') - \lambda(x))\} = \lambda(x) + \delta_f[\lambda(x)] ,\tag{8}$$

$$\hat{A}_\mu(x) = A_\mu(x) + \{\delta_f[A_\mu(x)] - (A_\mu(x') - A_\mu(x))\} = A_\mu(x) + A'_\mu(x) - A_\mu(x').\tag{9}$$

In the above we have defined,

$$\begin{aligned}x'_\mu &= x_\mu - f_\mu \quad , \quad A'_\mu(x') = \frac{\partial x^\nu}{\partial x'^\mu}A_\nu(x) \quad , \quad \lambda'(x') = \lambda(x) , \\ f^\mu &\equiv \frac{1}{2}\theta^{\mu\nu}A_\nu \quad .\end{aligned}\tag{10}$$

Here f^μ is the field dependent space-time translation parameter and δ_f constitutes the Lie derivative connected to f^μ ,

$$\begin{aligned}\delta_f[\lambda(x)] &= \lambda'(x) - \lambda(x) = -(\lambda(x') - \lambda(x)) = f^i\partial_i\lambda, \\ \delta_f[A_\mu(x)] &= A'_\mu(x) - A_\mu(x) .\end{aligned}\tag{11}$$

This shows that the NC gauge parameter ($\hat{\lambda}$) and gauge field ($\hat{A}^\mu$) are derivable from the ordinary one by making a *field dependent* space-time translation f^μ [21]. One can check that the NC gauge transformation of $\hat{A}_\mu(x)$ is correctly reproduced by considering,

$$\delta\hat{A}_\mu(x) = \delta(A_\mu(x) + \frac{1}{2}\theta^{\sigma\nu}A_\nu(x)F_{\sigma\mu}(x) + \frac{1}{2}\theta^{\sigma\nu}A_\nu(x)\partial_\sigma A_\mu(x)) ,\tag{12}$$

where $\delta A_\mu(x) = \partial_\mu \lambda(x)$ is the gauge transformation in ordinary space-time. Hence, if expressed in the form (9), the SWM, (at least to $O(\theta)$), can be derived in a geometrical way, without introducing the original identification (7) obtained from the viewpoint of a matching between NC and ordinary gauge invariant sectors. Also note that the gauge field $A_\mu(x)$ is treated here just as an ordinary vector field, without invoking any gauge theory properties. This constitutes the first part of our result.

Returning to our starting premises, are we justified in making an identification between $\hat{x}_\mu$ in (3)-(4) and x'_μ introduced in (8)-(10), because this relation can connect NC and ordinary space-time. Naively, a relation of the form, $x'_\mu = x_\mu - f_\mu(x)$ can not render the x' -space noncommutative, since the right hand side of the equation apparently comprises of commuting objects only. In our subsequent discussion we will show how this surmise can be made meaningful and will return to this point at the end.

IV. BATALIN-TYUTIN EXTENSION: $x_\mu \rightarrow \tilde{x}_\mu$ MAPPING

We start by considering a larger space having inherent NC. Such a space, which at the same time is physically appealing, is that of the Nambu-Goto model of relativistic spinning particle [19, 3]. Here the situation is somewhat akin to the open string boundary NC such that the role of Neveu-Schwarz field is played by here by the spin degrees of freedom. The Lagrangian of the model in 2+1-dimensions [19, 3] is,

$$L = [M^2 u^\mu u_\mu + \frac{J^2}{2} \sigma^{\mu\nu} \sigma_{\mu\nu} + M J \epsilon^{\mu\nu\lambda} u_\mu \sigma_{\nu\lambda}]^{\frac{1}{2}}, \quad (13)$$

$$u^\mu = \frac{dx^\mu}{d\tau}, \quad \sigma^{\mu\nu} = \Lambda_\rho{}^\mu \frac{d\Lambda^{\rho\nu}}{d\tau} = -\sigma^{\nu\mu}, \quad \Lambda_\rho{}^\mu \Lambda^{\rho\nu} = \Lambda^\mu{}_\rho \Lambda^{\nu\rho} = g^{\mu\nu}, \quad g^{00} = -g^{ii} = 1. \quad (14)$$

Here $(x^\mu, \Lambda^{\mu\nu})$ is a Poincare group element and also a set of dynamical variables of the theory.

In a Hamiltonian formulation, the conjugate momenta are,

$$P^\mu = \frac{\partial L}{\partial u_\mu} = L^{-1} [M^2 u^\mu + \frac{M J}{2} \epsilon^{\mu\nu\lambda} \sigma_{\nu\lambda}], \quad S^{\mu\nu} = \frac{\partial L}{\partial \sigma_{\mu\nu}} = \frac{L^{-1}}{2} [J^2 \sigma^{\mu\nu} + M J \epsilon^{\mu\nu\lambda} u_\lambda]. \quad (15)$$

The Poisson algebra of the above phase space degrees of freedom are,

$$\{P^\mu, x^\nu\} = g^{\mu\nu}, \quad \{P^\mu, P^\nu\} = 0, \quad \{x^\mu, x^\nu\} = 0, \quad \{\Lambda^{0\mu}, \Lambda^{0\nu}\} = 0, \quad (16)$$

$$\{S^{\mu\nu}, S^{\lambda\sigma}\} = S^{\mu\lambda} g^{\nu\sigma} - S^{\mu\sigma} g^{\nu\lambda} + S^{\nu\sigma} g^{\mu\lambda} - S^{\nu\lambda} g^{\mu\sigma}, \quad \{\Lambda^{0\mu}, S^{\nu\sigma}\} = \Lambda^{0\nu} g^{\mu\sigma} - \Lambda^{0\sigma} g^{\mu\nu}. \quad (17)$$

The full set of constraints are,

$$\Psi_1 \equiv P^\mu P_\mu - M^2 \approx 0, \quad \Psi_2 \equiv S^{\mu\nu} S_{\mu\nu} - 2J^2 \approx 0, \quad (18)$$

$$\Theta_1^\mu \equiv S^{\mu\nu} P_\nu, \quad \Theta_2^\mu \equiv \Lambda^{0\mu} - \frac{P^\mu}{M}, \quad \mu = 0, 1, 2, \quad (19)$$

out of which Ψ_1 and Ψ_2 give the mass and spin of the particle respectively.⁴ In the Dirac constraint analysis [17], these are termed as First Class Constraints (FCC), having the property that they commute with *all* the constraints on the constraint surface and generate gauge transformations. The set Θ_2^μ is put by hand [3], to restrict the number of angular co-ordinates.

The non-commuting set of constraints Θ_α^μ , $\alpha = 1, 2$, termed as Second Class Constraints (SCC) [17], modify the Poisson Brackets (16) to Dirac Brackets [17], defined below for any two generic variables A and B ,

$$\{A, B\}_{DB} = \{A, B\} - \{A, \Theta_\alpha^\mu\} \Delta_{\mu\nu}^{\alpha\beta} \{\Theta_\beta^\nu, B\}, \quad (20)$$

$$\{\Theta_\alpha^\mu, \Theta_\beta^\nu\} \equiv \Delta_{\alpha\beta}^{\mu\nu} \quad , \quad \alpha, \beta = 1, 2 \quad , \quad \Delta_{\alpha\beta}^{\mu\nu} \Delta_{\nu\lambda}^{\beta\gamma} = \delta_\alpha^\gamma \delta_\lambda^\mu . \quad (21)$$

$\Delta_{\alpha\beta}^{\mu\nu}$ is non-vanishing even on the constraint surface. The main result, relevant to us, is the following Dirac Bracket [19, 3],

$$\{x_\mu, x_\nu\}_{DB} = -\frac{S_{\mu\nu}}{M^2} \rightarrow \{\hat{x}_\mu, \hat{x}_\nu\} = \theta_{\mu\nu}. \quad (22)$$

This is the non-commutativity that occurs naturally in the spinning particle model. Our aim is to express this NC co-ordinate $\hat{x}_\mu$ in the form $\hat{x}_\mu = x_\mu - f_\mu$, with the identification between $\theta_{\mu\nu}$ and $S_{\mu\nu}$. This is indicated in the last equality in (22). In the quantum theory, this will lead to the NC space-time (3).

This motivates us to the Batalin-Tyutin quantization [18] of the spinning particle [3]. For a system of irreducible SCCs, in this formalism [18], the phase space is extended by introducing additional BT variables, ϕ_a^α , obeying

$$\{\phi_\mu^\alpha, \phi_\nu^\beta\} = \omega_{\mu\nu}^{\alpha\beta} = -\omega_{\nu\mu}^{\beta\alpha} \quad , \quad \omega_{\mu\nu}^{\alpha\beta} = g_{\mu\nu} \epsilon^{\alpha\beta} \quad , \quad \epsilon^{12} = 1. \quad (23)$$

where the last expression is a simple choice for $\omega_{\mu\nu}^{\alpha\beta}$. The SCCs Θ_α^μ are modified to $\tilde{\Theta}_\alpha^\mu$ such that they become FCC,

$$\{\tilde{\Theta}_\alpha^\mu(q, \phi), \tilde{\Theta}_\beta^\nu(q, \phi)\} = 0 \quad ; \quad \tilde{\Theta}_\alpha^\mu(q, \phi) = \Theta_\alpha^\mu(q) + \Sigma_{n=1}^\infty \tilde{\Theta}_\alpha^{\mu(n)}(q, \phi) \quad ; \quad \tilde{\Theta}^{\mu(n)} \approx O(\phi^n), \quad (24)$$

with q denoting the original degrees of freedom. Let us introduce the gauge invariant variables $\tilde{f}(q)$ [18] corresponding to each $f(q)$, so that $\{\tilde{f}(q), \tilde{\Theta}_\alpha^\mu\} = 0$

$$\tilde{f}(q, \phi) \equiv f(\tilde{q}) = f(q) + \Sigma_{n=1}^\infty \tilde{f}(q, \phi)^{(n)}, \quad (25)$$

which further satisfy [18],

$$\{q_1, q_2\}_{DB} = q_3 \rightarrow \{\tilde{q}_1, \tilde{q}_2\} = \tilde{q}_3 \quad , \quad \tilde{0} = 0. \quad (26)$$

It is now clear that our target is to obtain $\tilde{x}_\mu$ for x_μ . Explicit expressions for $\tilde{\Theta}^{\mu(n)}$ and $\tilde{f}^{(n)}$ are derived in [18].

⁴Note that instead of Ψ_2 as above, one can equivalently use $\Psi_2 \equiv \epsilon^{\mu\nu\lambda} S_{\mu\nu} P_\lambda - M J$, which incidentally defines the Pauli Lubanski scalar.

Before we plunge into the BT analysis, the reducibility of the SCCs Θ_1^μ (*i.e.* $P_\mu \Theta_1^\mu = 0$) [19, 3] is to be removed [22] by introducing a canonical pair of auxiliary variables ϕ and π that satisfy $\{\phi, \pi\} = 1$ and PB commute with the rest of the physical variables. The modified SCCs that appear in the subsequent BT analysis are as shown below:

$$\Theta_1^\mu \equiv S^{\mu\nu} P_\nu + k_1 P^\mu \pi \quad ; \quad \Theta_2^\mu \equiv (\Lambda^{0\mu} - \frac{P^\mu}{M}) + k_2 (\Lambda^{0\mu} + \frac{P^\mu}{M}) \phi , \quad (27)$$

where k_1 and k_2 denote two arbitrary parameters. Since the computations are exhaustively done in [3] they are not repeated here. The results are the following:

$$\tilde{x}_\mu = x_\mu + [S_{\nu\mu} + 2k_1 \pi g_{\nu\mu}] (\phi^1)^\nu + \mathcal{R}_{1\mu\nu} (\phi^2)^\nu + \text{higher} - \phi - \text{terms} , \quad (28)$$

$$\{\tilde{x}_\mu, \tilde{x}_\nu\} = -\frac{\tilde{S}_{\mu\nu}}{M^2} , \quad \tilde{S}_{\mu\nu} = S_{\mu\nu} + \mathcal{R}_{2(\alpha)\mu\nu\lambda} \phi^{(\alpha)\lambda} + \text{higher} - \phi - \text{terms} , \quad (29)$$

where the expressions for $\mathcal{R}$ are straightforward to obtain [3] but are not needed in the present order of analysis. Only it should be remembered that the $\mathcal{R}_1$ -term in (28) is responsible for the $(\phi^\alpha)^\mu$ -free term $-S_{\mu\nu}/(M^2)$ in the $\{\tilde{x}_\mu, \tilde{x}_\nu\}$ bracket in (29). Thus the problem that we had set out to solve has been addressed successfully in (28), which expresses the NC $\tilde{x}_\mu$ in terms of ordinary x_μ and other variables [3].

V. BATALIN-TYUTIN EXTENSION IN A GAUGE $\rightarrow$ THE SEIBERG-WITTEN MAP

Now comes the crucial part of identification of the present map with the SWM [7]. This means in particular that we have to connect (28) to (10), since as we have shown before, (10) is capable of generating the SWM [7]. We exploit the freedom of choosing gauges according to our convenience, since in the BT extended space $\tilde{\Theta}_\alpha^\mu$ are FCCs. For instance, the so called unitary gauge, $\phi_1^\mu = 0, \phi_2^\mu = 0$, trivially converts the system back to its original form before the BT extension. Let us choose the following non-trivial gauge,

$$\phi_1^\mu = \frac{M^2}{2} A^\mu(x) \quad , \quad \phi_2^\mu = 0, \quad (30)$$

where $A^\mu(x)$ is some function of x_μ , to be identified with the gauge field. Let us also work with terms linear in $A^\mu(x)$. Identifying $\tilde{S}_{\mu\nu}/(M)^2 = \theta_{\mu\nu}$ we end up with the cherished mapping,

$$\tilde{x}_\mu = x_\mu - \frac{1}{2} \theta_{\mu\nu} A^\nu(x) + \text{higher} - A(x) - \text{terms} , \quad (31)$$

$$\{\tilde{x}_\mu, \tilde{x}_\nu\}_{DB} = \theta_{\mu\nu} + \text{higher} - A(x) - \text{terms} . \quad (32)$$

Note that in the above relations (31,32), we have dropped the terms containing k_1 , an arbitrary parameter [3], considering it to be very small. Also in (32) Dirac Bracket reappears since the system is gauge fixed and hence has SCCs. This constitutes the second part of our result.

Finally, two points are to be noted. Firstly, the non-commutativity present here does *not* break Lorentz invariance since there appear no constant parameter with non-trivial Lorentz index to start with. The violation will appear only in the identification of $\tilde{S}_{\mu\nu}$ with (constant) $\theta_{\mu\nu}$. Secondly, (28) truly expresses the NC space-time $\tilde{x}_\mu$ in terms of ordinary space-time x_μ . But x_μ becomes NC owing to the Dirac brackets induced by the particular gauge that we fixed in order to reduce our results to the SWM. Obviously, in general, there is no need to fix this particular gauge. This refers to the comment below (12).

VI. CONCLUSIONS AND FUTURE PROSPECTS

In conclusion, let us summarize our work. We have shown that the (abelian $O(\theta)$) Seiberg-Witten map can be viewed as a co-ordinate transformation, albeit with field dependent parameters. The duality concept between gauge orbits in noncommutative and ordinary space-times, which was crucial in the original derivation [7], is not applied here. It has been explicitly demonstrated that a noncommutative space-time sector can be constructed in the Batalin-Tyutin extension of the relativistic spinning particle model [3]. Finally, the above mentioned transformation and subsequently a direct connection with the Seiberg-Witten map is also generated in this model. The present work reveals that noncommutative space-time is endowed with spin degrees of freedom, as compared to the ordinary space-time [5].

As a future work, we plan to develop the quantum theory of the noncommutative spacetime model proposed in [5] whose particle content is analogous to the spinning particles considered here. This theory has a classical conformal invariance, the fate of which will be studied upon quantization. Also, it would be interesting to investigate how the constraints induce the noncommutativity in spacetime coordinates in an operator product expansion approach.

Acknowledgement: It is a pleasure to thank Professor R.Jackiw for helpful correspondence.

References

- [1] H.S.Snyder, Phys.Rev. 71 38(1947).
- [2] S.Doplicher, K.Fredenhagen and J.E.Roberts, Phys.Lett. B331 39 (1994); C.E.Carlson, C.D.Carrone and N.Zobin, *Non-commutative gauge theory without Lorentz violation*, Phys.Rev. D66 (2002) 075001 (hep-th/0206035).
- [3] S.Ghosh, Phys.Rev. D66 045031(2002).
- [4] H.Kase, K.Morita, Y.Okumura and E.Umezawa, *Lorentz- Invariant Non-Commutative Space-Time Based On DFR Algebra*, hep-th/0212127.
- [5] S.Ghosh, *Modelling a noncommutative two-brane*, hep- th/0212123.
- [6] For a review see M.R.Douglas and N.A.Nekrasov, Rev.Mod.Phys. 73 977(2002) (arXiv: HEP-TH/ 0106048); R.J.Szabo, hep-th/0109162; A.Connes, *Noncommutative Geometry*, Academic Press, 1994.
- [7] N.Seiberg and E.Witten, JHEP 09(1999)032.
- [8] R.Banerjee, B.Chakraborty and S.Ghosh, Phys.Lett. B537 340 (2002).
- [9] Y.Okawa and H.Ooguri, Phys.Rev. D64 046009(2001); S.Fidanza, JHEP 0206 016(2002).
- [10] X.Calmet et. al., hep-ph/0111115; H. Bozkaya, P. Fischer, H. Grosse, M. Pitschmann, V. Putz, M. Schweda, R. Wulkenhaar, *Space/time noncommutative field theories and causality*, HEP- TH/0209253; M.Chaichian, K.Nishijima and A.Tureanu, *Spin-Statistics and CPT Theorems in Noncommutative Field Theory*, hep-th/0209008; T.Tamaki, T.Harada, U.Miyamoto and T.Torii, *Particle velocity in noncommutative space-time*, Phys.Rev. D66 (2002) 105003 (gr-qc/0208002); J.L.Cortes, J.Gamboa, F.Mendez, *Noncommutativity in Field Space and Lorentz Invariance Violation*, J.M.Carmona, hep-th/0207158.
- [11] R.Banerjee and S.Ghosh, Phys.Lett. B533 162(2002).
- [12] J.M.Gracia-Bondia and C.P.Martin, Phys.Lett. B479 321(2000); L.Bonora, M.Schnabl and A.Tomasiello, Phys.Lett. B485 311(2000); F.Ardalan and N.Sadooghi, Int.J.Mod.Phys. A16 3157(2001).
- [13] S.Ghosh, Phys.Lett. B 558 245 (2003) (hep-th/0210107); M.B.Cantcheff and P.Minces, Phys.Lett. B557 283(2003) (hep-th/0212031); O.F.Dayi, Phys.Lett. B560 239(2003).
- [14] S.Deser, R.Jackiw and S.Templeton, Phys.Rev.Lett. 48(1982)975; Ann.Phys. 140 (1982)372; S.Deser and R.Jackiw, Phys.Lett. 139B(1984)371.
- [15] E.Fradkin and F.A.Schaposnik, Phys.Lett. 338B(1994)253; R.Banerjee, Phys.Lett. 358B (1995)297.

- [16] S.Ghosh, Phys.Lett. B563 112(2003).
- [17] P.A.M.Dirac, *Lectures on Quantum Mechanics* (Yeshiva University Press, New York, 1964).
- [18] I.A.Batalin and I.V.Tyutin, Int.J.Mod.Phys. A6 3255(1991).
- [19] S.Ghosh, Phys.Lett. B338 235(1994); J.Math.Phys. 42 5202 (2001). For the original works see, A.J.Hanson and T.Regge, Ann.Phys. (N.Y.) 87 498(1974); see also A.J.Hanson, T.Regge and C.Teitelboim, *Constrained Hamiltonian System*, Roma, Accademia Nazionale Dei Lincei, (1976).
- [20] L.Cornalba, *D-brane physics and noncommutative Yang-Mills theory*, hep-th/9909081; A.A.Bichl et. al., *Noncommutative Lorentz symmetry and the origin of the Seiberg-Witten map*, hep-th/0108045; B.Jurco and P.Schupp, Euro.Phys.J. C14 367(2000); B.Jurco and P.Schupp and J.Wess, Nucl.Phys. B584 784(2001); H.Liu, Nucl.Phys. B614 305(2001); R.Jackiw, S.-Y.Pi and A.P.Polychronakos, *Noncommutating gauge fields as a Lagrange fluid*, Phys.Rev.Lett. 88 111603(2002) (hep-th/0206014).
- [21] Field dependent transformations have appeared in *eg* R.Jackiw, Phys.Rev.Lett. 41, 1635 (1978); R. Jackiw and S.-Y. Pi, hep-th/0111122.
- [22] R.Banerjee and J.Barcelos-Neto, Ann.Phys. 265 134(1998).